

ON BURKHOLDER FUNCTION FOR ORTHOGONAL MARTINGALES AND ZEROS OF LEGENDRE POLYNOMIALS

ALEXANDER BORICHEV, PRABHU JANAKIRAMAN, ALEXANDER VOLBERG

ABSTRACT. Burkholder obtained a sharp estimate of $\mathbf{E}|W|^p$ via $\mathbf{E}|Z|^p$, for martingales W differentially subordinated to martingales Z . His result is that $\mathbf{E}|W|^p \leq (p^* - 1)^p \mathbf{E}|Z|^p$, where $p^* = \max(p, \frac{p}{p-1})$. What happens if the martingales have an extra property of being orthogonal martingales? This property is an analog (for martingales) of the Cauchy-Riemann equation for functions, and it naturally appears in a problem on singular integrals (see the references at the end of Section 1). We establish here that in this case the constant is quite different. Actually, $\mathbf{E}|W|^p \leq (\frac{1+z_p}{1-z_p})^p \mathbf{E}|Z|^p$, $p \geq 2$, where z_p is a specific zero of a certain solution of the Legendre ODE. We also prove the sharpness of this estimate. Asymptotically, $(1+z_p)/(1-z_p) = (4j_0^{-2} + o(1))p$, $p \rightarrow \infty$, where j_0 is the first positive zero of the Bessel function of zero order. This connection with zeros of special functions (and orthogonal polynomials for $p = n(n+1)$) is rather unexpected.

1. INTRODUCTION

Let $Z = (X, Y)$, $W = (U, V)$ be two $\mathbb{R}^2$ -valued martingales on the filtration of the 2-dimensional Brownian motion $B_s = (B_{1s}, B_{2s})^T$. Let

$$A = \begin{bmatrix} -1, & i \\ i, & 1 \end{bmatrix}.$$

The research of the first author was partially supported by the ANR grants DYNOP and FRAB; the research of the second and the third authors was partially supported by the NSF grants DMS-0501067 and DMS-0605166.

We want W to be a martingale transform of Z defined by A . Let

$$\begin{aligned} X(t) &= \int_0^t \vec{x}(s) \cdot dB_s, \\ Y(t) &= \int_0^t \vec{y}(s) \cdot dB_s, \end{aligned}$$

where X, Y are *real-valued* processes, and $\vec{x}(s), \vec{y}(s)$ are $\mathbb{R}^2$ -valued “martingale differences” written as row vectors.

Put

$$Z(t) = X(t) + iY(t), \quad Z(t) = \int_0^t (\vec{x}(s) + i\vec{y}(s)) \cdot dB_s,$$

and

$$W(t) = U(t) + iV(t), \quad W(t) = \int_0^t (A(\vec{x}(s)^T + i\vec{y}(s)^T))^T \cdot dB_s.$$

We denote

$$W = A \star Z.$$

As above,

$$\begin{aligned} U(t) &= \int_0^t \vec{u}(s) \cdot dB_s, \\ V(t) &= \int_0^t \vec{v}(s) \cdot dB_s, \\ W(t) &= \int_0^t (\vec{u}(s) + i\vec{v}(s)) \cdot dB_s. \end{aligned}$$

We can easily write the components of $\vec{u}(s), \vec{v}(s)$:

$$\begin{aligned} u_1(s) &= -x_1(s) - y_2(s), & v_1(s) &= x_2(s) - y_1(s), \\ u_2(s) &= x_2(s) - y_1(s), & v_2(s) &= x_1(s) + y_2(s). \end{aligned}$$

Note that

$$\vec{u} \cdot \vec{v} = u_1v_1 + u_2v_2 = -(x_1 + y_2)(x_2 - y_1) + (x_2 - y_1)(x_1 + y_2) = 0.$$

1.1. Local orthogonality. The processes

$$\begin{aligned}
\langle X, U \rangle(t) &:= \int_0^t \vec{x} \cdot \vec{u} ds, & \langle X, V \rangle(t) &:= \int_0^t \vec{x} \cdot \vec{v} ds, \\
\langle Y, U \rangle(t) &:= \int_0^t \vec{y} \cdot \vec{u} ds, & \langle Y, V \rangle(t) &:= \int_0^t \vec{y} \cdot \vec{v} ds, \\
\langle X, X \rangle(t) &:= \int_0^t \vec{x} \cdot \vec{x} ds, & \langle Y, Y \rangle(t) &:= \int_0^t \vec{y} \cdot \vec{y} ds, \\
\langle X, Y \rangle(t) &:= \int_0^t \vec{x} \cdot \vec{y} ds, & \langle U, U \rangle(t) &:= \int_0^t \vec{u} \cdot \vec{u} ds, \\
\langle V, V \rangle(t) &:= \int_0^t \vec{v} \cdot \vec{v} ds, & \langle U, V \rangle(t) &:= \int_0^t \vec{u} \cdot \vec{v} ds.
\end{aligned}$$

are called the covariance processes. We can denote

$$\begin{aligned}
d\langle X, U \rangle(t) &:= \vec{x}(t) \cdot \vec{u}(t), & d\langle X, V \rangle(t) &:= \vec{x}(t) \cdot \vec{v}(t), \\
d\langle Y, U \rangle(t) &:= \vec{y}(t) \cdot \vec{u}(t), & d\langle Y, V \rangle(t) &:= \vec{y}(t) \cdot \vec{v}(t), \\
d\langle X, X \rangle(t) &:= \vec{x}(t) \cdot \vec{x}(t), & d\langle Y, Y \rangle(t) &:= \vec{y}(t) \cdot \vec{y}(t), \\
d\langle X, Y \rangle(t) &:= \vec{x}(t) \cdot \vec{y}(t), & d\langle U, U \rangle(t) &:= \vec{u}(t) \cdot \vec{u}(t), \\
d\langle V, V \rangle(t) &:= \vec{v}(t) \cdot \vec{v}(t), & d\langle U, V \rangle(t) &:= \vec{u}(t) \cdot \vec{v}(t), \\
d\langle Z, Z \rangle(t) &:= \vec{x}(t) \cdot \vec{x}(t) + \vec{y}(t) \cdot \vec{y}(t), \\
d\langle W, W \rangle(t) &:= \vec{u}(t) \cdot \vec{u}(t) + \vec{v}(t) \cdot \vec{v}(t).
\end{aligned}$$

The following observations are important.

Lemma 1. *Let $W = U + iV$ be the martingale transform of $Z = X + iY$ by means of the matrix A above. Then*

$$d\langle U, V \rangle(t) = 0, \quad d\langle U, U \rangle(t) = d\langle V, V \rangle(t). \quad (1.1)$$

Equivalently,

$$\vec{u}(t) \cdot \vec{v}(t) = 0, \quad |\vec{u}(t)| = |\vec{v}(t)|.$$

Lemma 2. *We have*

$$d\langle U, U \rangle(t) \leq 2 d\langle Z, Z \rangle(t), \quad d\langle V, V \rangle(t) \leq 2 d\langle Z, Z \rangle(t).$$

Equivalently,

$$\begin{aligned}\vec{u}(t) \cdot \vec{u}(t) &\leq 2(\vec{x}(t) \cdot \vec{x}(t) + \vec{y}(t) \cdot \vec{y}(t)), \\ \vec{v}(t) \cdot \vec{v}(t) &\leq 2(\vec{x}(t) \cdot \vec{x}(t) + \vec{y}(t) \cdot \vec{y}(t)).\end{aligned}$$

or, equivalently,

$$d\langle W, W \rangle(t) \leq 4 d\langle Z, Z \rangle(t).$$

Proof. We have

$$\vec{u}(t) \cdot \vec{u}(t) = (x_1 + y_2)^2 + (x_2 - y_1)^2 \leq 2(x_1^2 + y_2^2 + x_2^2 + y_1^2) = 2 d\langle Z, Z \rangle.$$

Similar calculations can be made for v . $\square$

Definition 3. The complex martingale $W = A \star Z$ will be called the Ahlfors–Beurling transform of the martingale Z .

Now let us quote a theorem of Banuelos–Janakiraman [2]:

Theorem 4. *Let Z, W be two martingales on the filtration of the 2-dimensional Brownian motion, and let W be an orthogonal martingale in the sense of (1.1): $d\langle U, V \rangle(t) = 0$, $d\langle U, U \rangle(t) = d\langle V, V \rangle(t)$. Suppose that Z and W satisfy the subordination property*

$$d\langle W, W \rangle \leq d\langle Z, Z \rangle.$$

Let $p \geq 2$. Then for every t ,

$$(\mathbf{E} |W(t)|^p)^{1/p} \leq \sqrt{\frac{p^2 - p}{2}} (\mathbf{E} |Z(t)|^p)^{1/p}$$

($|\cdot|$ denotes the euclidean norm in $\mathbb{R}^2$).

One can easily obtain a “dual” version for $1 < p \leq 2$:

Theorem 5. *Let Z, W be two martingales on the filtration of the 2-dimensional Brownian motion, and let Z be an orthogonal martingale in the sense of (1.1): $d\langle X, Y \rangle(t) = 0$, $d\langle X, X \rangle(t) = d\langle Y, Y \rangle(t)$. Suppose that Z and W satisfy the subordination property*

$$d\langle W, W \rangle \leq d\langle Z, Z \rangle.$$

Let $1 < p \leq 2$. Then for every t ,

$$(\mathbf{E} |W(t)|^p)^{1/p} \leq \sqrt{\frac{2}{p^2 - p}} (\mathbf{E} |Z(t)|^p)^{1/p}.$$

We use the notations

$$\|Z(t)\|_p := (\mathbf{E} |Z(t)|^p)^{1/p}.$$

Sometimes we omit t and just write $\|Z\|_p$.

Theorem 4 together with Lemmas 1, 2 gives the following statement.

Theorem 6. $\|W\|_p = \|A \star Z\|_p \leq \sqrt{2(p^2 - p)} \|Z\|_p$, $p \geq 2$.

What happens in Theorems 4 for $1 < p < 2$ and in Theorem 5 for $p > 2$ is quite interesting, especially because these problems have such a close connection to estimates of the Ahlfors–Beurling operator, and because these problems exercise a lot of resistance. See also [5] and [37, Section 5] where it is shown how a big class of singular operators can be obtained from martingale transforms.

Acknowledgments. We are very grateful to Vasily Vasyunin who read the preliminary text and made many useful remarks. We are also very grateful to Fedja Nazarov for useful discussion and valuable remarks.

2. LEFT-RIGHT ORTHOGONALITY. MAIN THEOREM

As above, let us consider two $\mathbb{R}^2$ -valued martingales $Z = (X, Y)$ and $W = (U, V)$, both on the filtration of the 2-dimensional Brownian motion. Using the previous notations, we write

$$\vec{x}(t) = \nabla X(t), \quad \vec{y}(t) = \nabla Y(t), \quad \vec{u}(t) = \nabla U(t), \quad \vec{v}(t) = \nabla V(t).$$

Here the symbol ∇ stands for “stochastic gradient” of our martingales, somewhat abusing the notations. We assume the pointwise orthogonality

$$\vec{x}(t) \cdot \vec{y}(t) = 0, \quad |\vec{x}(t)| = |\vec{y}(t)|, \quad \vec{u}(t) \cdot \vec{v}(t) = 0, \quad |\vec{u}(t)| = |\vec{v}(t)|. \quad (2.1)$$

We also assume the pointwise subordination

$$|\vec{u}(t)|^2 + |\vec{v}(t)|^2 \leq |\vec{x}(t)|^2 + |\vec{y}(t)|^2. \quad (2.2)$$

To formulate our main result, let us recall that the Legendre function $L_\alpha(s)$ of order α is the unique (up to a multiplicative constant) bounded near 1 solution of the Sturm–Liouville equation

$$((1-s^2)y'(s))' + \alpha(\alpha+1)y(s) = 0.$$

It is the hypergeometric function ${}_2F_1(-\alpha, \alpha+1, 1; \frac{1-s}{2})$.

For any $p \in (2, \infty)$ consider the positive number α_p such that $\alpha_p(\alpha_p+1) = p$ that is

$$\alpha_p = \frac{\sqrt{1+4p}-1}{2}.$$

Consider the **largest** zero of $L_{\alpha_p}(s)$ on $(0, 1)$. Denote it by z_p .

Theorem 7. *Let $p \geq 2$, let Z, W satisfy (2.1) and (2.2). Then*

$$\|W\|_p \leq \frac{1+z_p}{1-z_p} \|Z\|_p.$$

Moreover, this constant is sharp.

Remark. In a recent preprint of Bañuelos and Osekowski [4] this theorem is extended to $0 < p < 2$. Moreover, it is extended to conformal martingales in $\mathbb{R}^d$. The extension is in the language of Bessel processes, which allows for non-integer d as well!

Let J_0 be the Bessel function of the zero order,

$$J_0(x) = \sum_{n \geq 0} (-1)^n \frac{x^{2n}}{2^{2n}(n!)^2}.$$

Denote its first positive zero by j_0 . It is known (see, for example, [39, Section 15.51]) that

$$j_0 \approx 2.4048.$$

Furthermore (see Section 10),

$$\lim_{p \rightarrow \infty} \frac{1}{p} \cdot \frac{1+z_p}{1-z_p} = \frac{4}{j_0^2}.$$

Since $j_0 > 2\sqrt[4]{2}$, our theorem gives better linear asymptotics for the constant as $p \rightarrow \infty$ than that in Theorem 4.

To prove Theorem 7 we are going to introduce the following Bellman function, a variant of Burkholder's function from [7]–[10], which will work for orthogonal martingales.

3. THE BELLMAN FUNCTION

We consider the martingales given by the stochastic integrals

$$\begin{aligned} X(t) &= X(0) + \int_0^t \vec{x}(s) \cdot dB_s, & Y(t) &= Y(0) + \int_0^t \vec{y}(s) \cdot dB_s, \\ U(t) &= U(0) + \int_0^t \vec{u}(s) \cdot dB_s, & V(t) &= V(0) + \int_0^t \vec{v}(s) \cdot dB_s, \\ M(t) &= M(0) + \int_0^t \vec{m}(s) \cdot dB_s. \end{aligned}$$

We assume that the martingale $Z(t) = (X(t), Y(t))$ satisfies the condition

$$\lim_{t \rightarrow \infty} \mathbf{E} |Z(t)|^p < \infty.$$

(Since $\mathbf{E} |Z(t)|^p$ is monotone, we have $\lim_{t \rightarrow \infty} \mathbf{E} |Z(t)|^p = \sup_{t \geq 0} \mathbf{E} |Z(t)|^p$.)

The martingale $M(t)$ is any martingale majorizing the sub-martingale $|Z(t)|^p$.

Here we assume that the 5×2 -matrix of random processes

$$e(s) := (\vec{x}(s), \vec{y}(s), \vec{u}(s), \vec{v}(s), \vec{m}(s))^T$$

satisfies pointwise the condition $e(s) \in A$, where

$$\begin{aligned} A = \{e \in M_{5 \times 2} : e_{11}e_{21} + e_{12}e_{22} = 0, e_{11}^2 + e_{12}^2 = e_{21}^2 + e_{22}^2, e_{31}e_{41} + e_{32}e_{42} = 0, \\ e_{31}^2 + e_{32}^2 = e_{41}^2 + e_{42}^2, e_{31}^2 + e_{32}^2 + e_{41}^2 + e_{42}^2 \leq e_{11}^2 + e_{12}^2 + e_{21}^2 + e_{22}^2\}. \end{aligned}$$

The random process $e(s)$ is also assumed to be a non-anticipatory process, that is, $e(s)$ is measurable with respect to the σ -algebra generated by the 2-dimensional Brownian motion $\{B_\tau, \tau \leq s\}$.

Introduce now $W = (U, V)$. For $T = (T_1, T_2, T_3, T_4, T_5) \in \mathbb{R}^5$ we define

$$\begin{aligned} \mathcal{B}(T) := & \sup_{\text{all non-anticipatory processes } e_{ij} \text{ such that } e(s) \in A,} \{ \lim_{t \rightarrow \infty} \mathbf{E} |W(t)|^p \}. \\ & X(0) = T_1, Y(0) = T_2, U(0) = T_3, V(0) = T_4, M(0) = T_5, M(t) \geq |Z(t)|^p \end{aligned}$$

It is convenient to change the notations and write everything in the following more compact form of a motion in $\mathbb{R}^5$:

$$R(t) = R(0) + \int_0^t e(s) dB_s.$$

Then

$$\mathcal{B}(T) := \sup_{\substack{\text{all non-anticipatory processes } e_{ij} \text{ such that } e(s) \in A, \\ R(0) = T, R_5(t) \geq (R_1(t)^2 + R_2(t)^2)^{p/2}}} \left\{ \lim_{t \rightarrow \infty} \mathbf{E} (R_3(t)^2 + R_4(t)^2)^{p/2} \right\}.$$

The function $\mathcal{B}(T)$ is defined on the following convex domain inside $\mathbb{R}^5$:

$$\Omega := \{T \in \mathbb{R}^5 : T_5 \geq (T_1^2 + T_2^2)^{p/2}\}.$$

3.1. Properties of the Bellman function. Let us fix a positive time t . Choose any non-anticipatory process $e(\tau)$, $0 \leq \tau < t$, satisfying the above restrictions. If we start from a point $R(0) \in \Omega$, we obtain the points $P_t = R(t, \omega)$. These are our starting data now.

Choose a matrix process $e(s)$, $s \geq t$, for a given $R(t, \omega)$, that attains the supremum in the definition of $\mathcal{B}(R(T))$ up to a small $\varepsilon > 0$. Then the process e , equal to $e(\tau)$, $0 \leq \tau < t$, $e(s)$, $s > t$, should be compared to the processes for the starting data $T = R(0)$ giving the supremum in the definition of $\mathcal{B}(R(0))$. Let us do this comparison. Introduce

$$F(T) = (T_3^2 + T_4^2)^{p/2}.$$

Let $R(t)$ be the martingale driven by e constructed above. Using the formula of full probability and stationarity of Brownian motion B_s we can write “Bellman’s principle”:

$$\mathcal{B}(R(0)) \geq \mathbf{E} F(R(\infty)) = \mathbf{E} \mathbf{E} (F(R(\infty)) | R(t, \omega) = P_t) \geq \mathbf{E} \mathcal{B}(R(t)) - \varepsilon.$$

In other words,

$$\mathbf{E} (\mathcal{B}(R(t)) - \mathcal{B}(R(0))) \leq 0.$$

Since F is convex, and $R(t)$ satisfies the martingale property for any initial point $T = R(0)$ in Ω , we have obviously

$$\mathcal{B}(T) \geq \mathcal{B}(R(t)) \geq \mathbf{E} F(R(\infty)) \geq F(\mathbf{E} R(\infty)) = F(R(0)) = (T_3^2 + T_4^2)^{\frac{p}{2}}.$$

Now we apply the Itô formula for the difference:

$$\mathbf{E}(\mathcal{B}(R(t)) - \mathcal{B}(R(0))) = \frac{1}{2} \int_0^t \mathbf{E} \sum_{k,l=1}^5 \frac{\partial^2 \mathcal{B}}{\partial T_k \partial T_l} (dR_k(s) \cdot dR_l(s)) ds;$$

we use here that $\mathbf{E}|B_{s+\Delta s} - B_s|^2 = \Delta s$. Note that by the formulas at the beginning of the section, $dR_j(s)$ is exactly $e_j(s)$, that is the j -th row of the matrix e . Therefore,

$$\begin{aligned} \mathbf{E}(\mathcal{B}(R(t)) - \mathcal{B}(R(0))) &= \frac{1}{2} \int_0^t \mathbf{E} \sum_{k,l=1}^5 \frac{\partial^2 \mathcal{B}}{\partial T_k \partial T_l} (\vec{e}_k(s) \cdot \vec{e}_l(s)) ds \\ &= \frac{1}{2} \int_0^t \mathbf{E} \text{trace}(e(s)^T d^2 \mathcal{B} e(s)) ds \leq 0, \end{aligned}$$

where $d^2 \mathcal{B}$ denotes the Hessian (matrix) of $\mathcal{B}$.

This formula holds for all non-anticipatory matrix processes $e(s)$, $0 \leq s \leq t$. We have already tacitly assumed the smoothness of $\mathcal{B}$. Using this assumption again, we divide the latter inequality by t and pass to the limit $t \rightarrow 0$. Then we get:

$$-\text{trace}(e(s)^T d^2 \mathcal{B} e(s)) = - \sum_{k,l=1}^5 \frac{\partial^2 \mathcal{B}}{\partial T_k \partial T_l} (\vec{e}_k \cdot \vec{e}_l) \geq 0 \quad e \in A. \quad (3.1)$$

Actually we might hope to have more (and these hopes will be, although only partially, fulfilled):

$$\max_{e \in A, e \neq 0} \sum_{k,l=1}^5 \frac{\partial^2 \mathcal{B}}{\partial T_k \partial T_l} (\vec{e}_k \cdot \vec{e}_l) = 0, . \quad (3.2)$$

We will not use (3.2) in the future. As we told, it was just a hope.

A more rigorous analysis of how to obtain (3.2) can be found in [21]. Still, it is not totally clear what conditions guarantee that “each state has the best control”.

Note that each vector e_j , $j = 1, \dots, 5$ has two coordinates. Let us unite all first coordinates and call the corresponding 5-vector e^1 , similarly we get e^2 . Then (3.1) can be rewritten as

$$\text{trace}(e^T d^2 \mathcal{B} e) = (d^2 \mathcal{B} e^1) \cdot e^1 + (d^2 \mathcal{B} e^2) \cdot e^2 \leq 0, \quad e \in A.$$

Similarly, we might hope to have

$$\max_{e \in A, e \neq 0} \text{trace}(e^T d^2 \mathcal{B} e) = \sup_{e=(e^1, e^2) \in A} ((d^2 \mathcal{B} e^1) \cdot e^1 + (d^2 \mathcal{B} e^2) \cdot e^2) = 0.$$

Theorem 8. *Let Ψ be any function such that*

$$\text{trace}(e(s)^T d^2 \Psi e(s)) \leq 0, \quad e \in A \quad (3.3)$$

(in the sense of distributions),

$$\Psi(T) \geq F(T), \quad (3.4)$$

and let for some $c > 0$ we have

$$\Psi(T) \leq c^p T_5, \quad (3.5)$$

for all T satisfying $T_1 = T_2 = T_3 = T_4 = 0$.

Then $\|W(t)\|_p \leq c\|Z(t)\|_p$, for any time t and any two orthogonal martingales W, Z such that W is differentially subordinated to Z in the sense of (2.2).

Proof. Fix $t \geq 0$ and denote $R_1 := X$, $R_2 := Y$, $R_3 := U$, $R_4 := V$, $M := |(R_1(t), R_2(t))|^p$. Consider the martingale $R_5(s) = \mathbf{E}(M|\mathcal{F}_s)$, where $\mathcal{F}_s$ is the σ -algebra generated by $B_1(\tau), B_2(\tau)$ for $\tau \leq s$. Let us consider $\mathbf{E} \Psi(R(s))$. If we apply Itô's formula to $\Psi(R(s))$ on $[0, t]$ and take the expectation, we get

$$\mathbf{E} \Psi(R(t)) = \frac{1}{2} \mathbf{E} \int_0^t \sum_{k,l=1}^5 \frac{\partial^2 \Psi}{\partial T_k \partial T_l} (dR_k(s) \cdot dR_l(s)) ds + \Psi(R(0)).$$

Next we use (3.3)–(3.5) and the convention $R_3 = U$, $R_4 = V$ to get

$$\mathbf{E} |(U(t), V(t))|^p \leq \Psi(R(0)) \leq c^p R_5(0).$$

Since $R_5(s)$ is a martingale, we have

$$R_5(0) = \mathbf{E} R_5(0) = \mathbf{E} R_5(t) = \mathbf{E} M = \mathbf{E} |(R_1(t), R_2(t))|^p.$$

It remains to note that we have the convention: $R_1 = X$, $R_2 = Y$. □

Corollary 9. *The best constant c such that $\mathcal{B}(T) \leq c^p T_5$ for all T satisfying $T_1 = T_2 = T_3 = T_4 = 0$ coincides with the best constant c such that $\|W(t)\|_p \leq c\|Z(t)\|_p$, for any time t and any two orthogonal martingales W, Z such that W is differentially subordinated to Z .*

It is easy to see several other properties of $\mathcal{B}$. For example, by multiplying our martingales T_1, T_2, T_3, T_4 by a positive constant τ one obtains:

$$\mathcal{B}(\tau T_1, \tau T_2, \tau T_3, \tau T_4, \tau^p T_5) = \tau^p \mathcal{B}(T).$$

Let S be a unitary operator on $\mathbb{R}^2$. Note that if we multiply its matrix by a matrix

$$\begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix}$$

such that its rows are orthogonal and the norms of rows are equal, then we have again a matrix with orthogonal rows having the same norm.

For us this means that given $(X, Y)^T, (U, V)^T$ (or $(R_1, R_2)^T, (R_3, R_4)^T$ in other notations), we can apply S to these vectors, and we can multiply the corresponding martingales for any t by the constant unitary matrix of S . Note that the process e will be transformed. Namely, the fifth row stays the same, but the rows 1, 2 and the rows 3, 4 form matrices which are multiplied on the left by the matrix of S . As we have found out, the new matrix e' has the same properties of rows as e ! So again $e' \in A$ pointwise. Of course, $|S \cdot (U, V)^T(t)| = |(U, V)^T(t)|$ pointwise. This reasoning gives us the following property of $\mathcal{B}$:

$$\mathcal{B}(T) =: b((T_1^2 + T_2^2)^{1/2}, (T_3^2 + T_4^2)^{1/2}, T_5).$$

The next property becomes obvious when we take $R_1(t) = R_3(t), R_2(t) = R_4(t)$:

$$(T_1^2 + T_2^2)^{p/2} \leq \mathcal{B}(T).$$

3.2. Reduction of the number of variables. Fix some $c > 0$ and $T' = (T_1, T_2, T_3, T_4)$. Recall that $(T', T_5) \in \Omega$ if and only if $T_5 \geq (T_1^2 + T_2^2)^{p/2}$, and consider

$$\Phi(T') = \Phi_c(T') = \sup_{T=(T', T_5): T \in \Omega} (\mathcal{B}(T) - c^p T_5).$$

If c is larger than the best constant in Corollary 9, then Φ is well defined at 0, and hence, everywhere.

Obviously, we have

$$\Phi(tT') = t^p \Phi(T'), \quad (3.6)$$

$$\Phi(T') =: \phi((T_1^2 + T_2^2)^{1/2}, (T_3^2 + T_4^2)^{1/2}). \quad (3.7)$$

By the submartingale property of $|W(t)|^p$ we obtain:

$$(T_3^2 + T_4^2)^{p/2} \leq \mathcal{B}(T).$$

What is much less easy (but still true) is that the concavity in the sense of (3.1) is also preserved. It is much less easy because the supremum of concave functions is not obliged to be concave. But if we have a concave function of several variables and form a new function which is the supremum of the original function over one of the variables, then the result is concave again. The same reasoning gives

$$-\text{trace}(e'^T d^2 \Phi e') = - \sum_{k,l=1}^4 \frac{\partial^2 \Phi}{\partial T_k \partial T_l} (\vec{e}_k \cdot \vec{e}_l) \geq 0, \quad e \in A. \quad (3.8)$$

Here e' denotes the matrix e with deleted fifth row.

Inequality (3.8) can be rewritten (again, in the sense of distributions) as

$$\text{trace}(e'^T d^2 \Phi e') = (d^2 \Phi e^1) \cdot e^1 + (d^2 \Phi e^2) \cdot e^2 \leq 0, \quad e \in \tilde{A}. \quad (3.9)$$

Here e^1, e^2 are 4-vectors, namely e^1 is the (column) vector of the first coordinates of all vectors $\vec{e}_k$, $k = 1, 2, 3, 4$, and e^2 is the (column) vector of the second coordinates of all vectors $\vec{e}_k$, $k = 1, 2, 3, 4$. We write $e^1 = (h, k)^T$, $e^2 = (h', k')^T$, where h, k, h', k' are (row) 2-vectors. Furthermore, the conditions on e' are the same as those on e but now with the fifth row deleted. We call these conditions $\tilde{A}$, and here they are:

$$|h| = |h'|, \quad |k| = |k'|, \quad h \cdot h' = 0, \quad k \cdot k' = 0, \quad |k| \leq |h|. \quad (3.10)$$

The next property is obvious:

$$(T_3^2 + T_4^2)^{p/2} - c^p (T_1^2 + T_2^2)^{p/2} \leq \Phi(T). \quad (3.11)$$

In the opposite direction, starting with a function Φ and $c > 0$ satisfying (3.6), (3.7), (3.9), (3.11), we can define

$$\Psi(T) = \Phi(T') + c^5 T_5,$$

and apply Theorem 8.

Corollary 10. *The best constant c such that there exists a function Φ satisfying (3.6), (3.7), (3.9), (3.11) coincides with the best constant c such that $\|W(t)\|_p \leq c\|Z(t)\|_p$, for any time t and any two orthogonal martingales W, Z such that W is differentially subordinated to Z .*

3.3. Another reduction of the number of variables. The function Φ has 4 variables, but the radial symmetry allows us to reduce it to a function of only 2 variables. Namely, using (3.6) and (3.7) we obtain a function $\phi(x, y)$, $x \geq 0, y \geq 0$ such that

$$\phi(tx, ty) = t^p \phi(x, y). \quad (3.12)$$

By (3.11), we have

$$y^p - c^p x^p \leq \phi(x, y).$$

Now we want to rewrite (3.9) in terms of ϕ using (3.10). Set

$$z = \frac{(T_1, T_2)}{|(T_1, T_2)|}, \quad w = \frac{(T_3, T_4)}{|(T_3, T_4)|}.$$

Given a vector h , we denote by $h^\perp$ the projection of h on the direction orthogonal to z ; given a vector k we denote by $k^\perp$ the projection of k on the direction orthogonal to w , the same with h', k' .

Set $x = (T_1^2 + T_2^2)^{1/2}$, $y = (T_3^2 + T_4^2)^{1/2}$, $u_1 = u_2 = x$, $u_3 = u_4 = y$. We have

$$\begin{aligned} \frac{\partial u_j}{\partial T_j} &= \frac{T_j}{u_j}, \quad \frac{\partial \Phi}{\partial T_j} = \frac{\partial \phi}{\partial u_j} \cdot \frac{T_j}{u_j}, \quad 1 \leq j \leq 4, \\ \frac{\partial^2 \Phi}{\partial T_j^2} &= \frac{\partial^2 \phi}{\partial u_j^2} \cdot \frac{T_j^2}{u_j^2} + \frac{\partial \phi}{\partial u_j} \cdot \frac{u_j^2 - T_j^2}{u_j^3}, \quad 1 \leq j \leq 4, \\ \frac{\partial^2 \Phi}{\partial T_j \partial T_k} &= \frac{\partial^2 \phi}{\partial u_j^2} \cdot \frac{T_j T_k}{u_j^2} - \frac{\partial \phi}{\partial u_j} \cdot \frac{T_j T_k}{u_j^3}, \quad 1 \leq j, k \leq 4, u_j = u_k, j \neq k, \\ \frac{\partial^2 \Phi}{\partial T_j \partial T_k} &= \frac{\partial^2 \phi}{\partial x \partial y} \cdot \frac{T_j T_k}{xy}, \quad 1 \leq j, k \leq 4, u_j \neq u_k. \end{aligned}$$

Then

$$\begin{aligned}
& (d^2\Phi(h, k)^T) \cdot (h, k)^T + (d^2\Phi(h', k')^T) \cdot (h', k')^T = \\
& \frac{1}{x} \frac{\partial \phi}{\partial x} |h^\perp|^2 + \frac{\partial^2 \phi}{\partial x^2} (h \cdot z)^2 + 2 \frac{\partial^2 \phi}{\partial x \partial y} (h \cdot z)(k \cdot w) + \frac{\partial^2 \phi}{\partial y^2} (k \cdot w)^2 + \frac{1}{y} \frac{\partial \phi}{\partial y} |k^\perp|^2 + \\
& \frac{1}{x} \frac{\partial \phi}{\partial x} |h'^\perp|^2 + \frac{\partial^2 \phi}{\partial x^2} (h' \cdot z)^2 + 2 \frac{\partial^2 \phi}{\partial x \partial y} (h' \cdot z)(k' \cdot w) + \frac{\partial^2 \phi}{\partial y^2} (k' \cdot w)^2 + \frac{1}{y} \frac{\partial \phi}{\partial y} |k'^\perp|^2 \\
& = \frac{1}{x} \frac{\partial \phi}{\partial x} |h|^2 + \frac{\partial^2 \phi}{\partial x^2} |h|^2 + 2 \frac{\partial^2 \phi}{\partial x \partial y} (h \cdot Sk) + \frac{\partial^2 \phi}{\partial y^2} |k|^2 + \frac{1}{y} \frac{\partial \phi}{\partial y} |k|^2.
\end{aligned}$$

Here S stands for a unitary operator sending w to z . Note that the last expression must be non-positive for any h, k such that

$$|k| \leq |h|.$$

Thus, (3.9) becomes (in the sense of distributions)

$$\frac{1}{x} \frac{\partial \phi}{\partial x} |h|^2 + \frac{\partial^2 \phi}{\partial x^2} |h|^2 + 2 \frac{\partial^2 \phi}{\partial x \partial y} (h \cdot k) + \frac{\partial^2 \phi}{\partial y^2} |k|^2 + \frac{1}{y} \frac{\partial \phi}{\partial y} |k|^2 \leq 0, \quad |k| \leq |h|. \quad (3.13)$$

4. REDUCTION TO DIFFERENTIAL INEQUALITIES IN ONE DIMENSION

We can reduce our problem to the following one. We are looking for the smallest c such that the function

$$h_c(x, y) := y^p - c^p x^p$$

can be majorized by a solution of the differential inequality (3.13) satisfying (3.12). Both ϕ and h_c are p -homogeneous. Therefore, we can further reduce our problem to that on functions of one real variable. First note that (3.13) is equivalent to the fact that a certain quadratic polynomial is negative when its argument is bigger than 1 in absolute value. This is equivalent to three inequalities, the first and the second of which are

$$\frac{1}{x} \frac{\partial \phi}{\partial x} + \frac{\partial^2 \phi}{\partial x^2} - 2 \frac{\partial^2 \phi}{\partial x \partial y} + \frac{\partial^2 \phi}{\partial y^2} + \frac{1}{y} \frac{\partial \phi}{\partial y} \leq 0, \quad (4.1)$$

$$\frac{1}{x} \frac{\partial \phi}{\partial x} + \frac{\partial^2 \phi}{\partial x^2} + 2 \frac{\partial^2 \phi}{\partial x \partial y} + \frac{\partial^2 \phi}{\partial y^2} + \frac{1}{y} \frac{\partial \phi}{\partial y} \leq 0. \quad (4.2)$$

These relations claim just that our quadratic expression is negative when its argument is equal to ± 1 .

Let $\mathcal{D}$ denote the discriminant of our quadratic expression,

$$\mathcal{D} = \left(\frac{\partial^2 \phi}{\partial x \partial y} \right)^2 - \left(\frac{1}{x} \frac{\partial \phi}{\partial x} + \frac{\partial^2 \phi}{\partial x^2} \right) \left(\frac{1}{y} \frac{\partial \phi}{\partial y} + \frac{\partial^2 \phi}{\partial y^2} \right).$$

Clearly, if $\mathcal{D} < 0$, and if the quadratic polynomial is negative at ± 1 by (4.1), (4.2), then it is negative for all the arguments exceeding 1 in absolute value and (3.13) holds. If $\mathcal{D} \geq 0$, then (3.13) follows from (4.1), (4.2), and the fact that the smaller root of the quadratic expression belongs to $(-1, 1)$, which is our third inequality:

$$\left| \left| \frac{\partial^2 \phi}{\partial x \partial y} \right| - \mathcal{D}^{1/2} \right| \leq \left| \frac{\partial^2 \phi}{\partial x^2} + \frac{1}{x} \frac{\partial \phi}{\partial x} \right|. \quad (4.3)$$

Now using homogeneity we write

$$s := \frac{y - x}{x + y}, \quad \frac{y}{x + y} = \frac{1 + s}{2}, \quad \frac{x}{x + y} = \frac{1 - s}{2}, \quad (4.4)$$

$$\begin{aligned} \phi(x, y) &= (x + y)^p \phi\left(\frac{x}{x + y}, \frac{y}{x + y}\right) = (x + y)^p \phi\left(\frac{1 - s}{2}, \frac{1 + s}{2}\right), \\ g(s) &:= \phi\left(\frac{1 - s}{2}, \frac{1 + s}{2}\right). \end{aligned}$$

On $\{(x, y) : x + y = 1\}$ we have:

$$\begin{aligned} \phi_{xx} &= p(p - 1)g(s) - 2(p - 1)(1 + s)g'(s) + (1 + s)^2 g''(s), \\ \phi_{yy} &= p(p - 1)g(s) + 2(p - 1)(1 - s)g'(s) + (1 - s)^2 g''(s), \\ \phi_{xy} &= p(p - 1)g(s) - 2(p - 1)sg'(s) - (1 - s^2)g''(s), \end{aligned} \quad (4.5)$$

$$\frac{\phi_x}{x} = \frac{2p}{1 - s}g(s) - \frac{2(1 + s)}{1 - s}g'(s), \quad (4.6)$$

$$\frac{\phi_y}{y} = \frac{2p}{1 + s}g(s) + \frac{2(1 - s)}{1 + s}g'(s), \quad (4.7)$$

$$\phi_{xx} - 2\phi_{xy} + \phi_{yy} = 4g''(s). \quad (4.8)$$

Denote

$$Kg(s) := p(p - 1)g(s) - 2(p - 1)sg'(s) - (1 - s^2)g''(s),$$

$$Dg(s) := ((1 - s^2)g'(s))' + pg(s),$$

$$\tilde{D}g(s) := \frac{Dg(s)}{1 - s^2} + Kg(s).$$

Then (4.1), (4.2) can be rewritten correspondingly as

$$Dg(s) = (1 - s^2)g''(s) - 2sg'(s) + pg(s) \leq 0, \quad s \in [-1, 1], \quad (4.9)$$

$$\tilde{D}g(s) = \frac{Dg(s)}{1 - s^2} + Kg(s) \leq 0, \quad s \in [-1, 1]. \quad (4.10)$$

Next we pass to (4.3). Suppose that $Dg = 0$ on an interval I . Then we can calculate on $\{(x, y) : x + y = 1, \frac{y-x}{x+y} \in I\}$:

$$\begin{aligned} \phi_{xy} &= p(-2sg' + pg), \\ \frac{1}{x}\phi_x + \phi_{xx} &= p(-2(1+s)g' + pg), \quad \frac{1}{y}\phi_y + \phi_{yy} = p(2(1-s)g' + pg). \end{aligned}$$

Therefore,

$$\mathcal{D} = 4p^2g'^2 \geq 0,$$

and condition (4.3) becomes

$$||-2sg' + pg| - 2|g'| \leq |-2(1+s)g' + pg|, \quad (4.11)$$

which is just the triangle inequality.

Thus, for any interval I where g is a solution of the Legendre equation $Dg = 0$, (4.3) is automatically satisfied on $\{(x, y) : \frac{y-x}{x+y} \in I\}$.

5. CASE $p = n(n+1)$

From now on, we assume that $p = \alpha(\alpha+1)$, $\alpha > 1$, and we use the notation D_α for the operator D defined in (4.9).

If $\alpha = n \in \mathbb{N}$, then $p = n(n+1)$ and the Legendre functions L_n (i.e. the solutions of the equation $D_n L_n = 0$ bounded near the point 1) become Legendre polynomials (see, for instance, [14, Section 3.8]):

$$L_n(s) = \frac{1}{2^n n!} \frac{d^n}{ds^n} (s^2 - 1)^n, \quad L_{n,a} = a L_n. \quad (5.1)$$

Note that $L_n(1) = 1$.

Let us consider an obstacle function

$$h_c(s) := \left(\frac{1+s}{2}\right)^p - c^p \left(\frac{1-s}{2}\right)^p.$$

Our first remark is that

$$D_\alpha h_c(s) = D_\alpha \left(\left(\frac{1+s}{2} \right)^p - c^p \left(\frac{1-s}{2} \right)^p \right) = \frac{p}{p-1} (1-s^2) h_c''(s).$$

In particular, the inflection point i_p of h_c coincides with the point where the function $D_\alpha h_c$ changes the sign from positive to negative (when we move from $s = 1$ to $s = -1$).

Suppose that $\alpha = n > 1$ is an integer. If $p = n(n+1)$, then the Legendre equation $D_n g = 0$ has two linearly independent solutions: one is the Legendre polynomial L_n of degree n , and another, Q_n , has logarithmic singularities at $x = \pm 1$.

The following statement is a partial case of Lemma 19 we prove later on:

Lemma 11. *Consider the set Y_p of all linear combinations of these solutions L_n and Q_n . For any $y \in Y_p$, let $f(y, p)$ be the largest zero of y on $[-1, 1]$. Then*

$$\min_{y \in Y_p} f(y, p) = f(L_n, p) =: z_p.$$

Consider all the pairs (f, h_c) , where f is a solution of the Legendre equation such that $f(1) > 1$ and h_c is as above (recall that $h_c(1) = 1$), $f > h_c$ on some interval $(x, 1]$, and f and h_c have the same values and the same derivatives at x . Here is a question which will occupy our attention almost till the end of this section:

Question: What is the smallest c possible for such pairs?

Theorem 12. *The smallest c is $\frac{1+z_p}{1-z_p}$.*

Consider the following situation: $L_{n,a}$ and h_c touch at a certain point $x = x(p, a)$, the first function being above the second one on $(x(p, a), 1]$. It is easy to obtain such a situation. Fix $a > 1$ to have $L_{n,a}(1) > 1$. Note that $h_c(1) = 1$ for every c . Take sufficiently large c . Since z_p is the first zero of L_n (counting from 1 to the left, $p = n(n+1)$), we have $L_{n,a} > h_c$ on $[z_p, 1]$. Start to decrease c . At a touching point $x = x(p, a)$, $c = c(p, a)$ we have the

equations

$$\begin{cases} \left(\frac{1+x}{2}\right)^p - c^p \left(\frac{1-x}{2}\right)^p = aL_n(x), \\ \frac{p}{2} \left(\left(\frac{1+x}{2}\right)^{p-1} + c^p \left(\frac{1-x}{2}\right)^{p-1}\right) = aL'_n(x). \end{cases}$$

Then

$$\begin{aligned} \left(\frac{1+x}{2}\right)^{p-1} &= a \left(\frac{1}{p}(1-x)L'_n(x) + L_n(x)\right), \\ c^p \left(\frac{1-x}{2}\right)^{p-1} &= a \left(\frac{1}{p}(1+x)L'_n(x) - L_n(x)\right), \end{aligned}$$

and we get

$$c^p = \left(\frac{1+x}{1-x}\right)^{p-1} \frac{\frac{1}{p}(1+x)L'_n(x) - L_n(x)}{\frac{1}{p}(1-x)L'_n(x) + L_n(x)}.$$

Let us consider the function

$$\beta(x) := \frac{(1+x)^p L'_n(x) - p(1+x)^{p-1} L_n(x)}{(1-x)^p L'_n(x) + p(1-x)^{p-1} L_n(x)}.$$

Lemma 13. *The function β is strictly increasing on the interval $[z_p, 1)$.*

Proof. Let us differentiate β and use the fact that L_n satisfies the equation $(1-x^2)L''_n(x) - 2xL'_n(x) + pL_n(x) = 0$. Then we get

$$\begin{aligned} & \frac{\beta'(x)((1-x)^p L'_n(x) + p(1-x)^{p-1} L_n(x))^2}{2p(1-x^2)^{p-2}} \\ &= L_n(x)((1-x^2)L''_n(x) + (p-1) \cdot 2xL'_n(x) - p(p-1)L_n(x)) \\ &= p(1-x^2) L_n(x) L''_n(x). \end{aligned} \tag{5.2}$$

The only zero of L_n on $[z_p, 1)$ is z_p . The polynomial L_n is positive on $(z_p, 1)$. Let us show that

$$L''_n(s) > 0, \quad s \in [z_p, 1). \tag{5.3}$$

The orthogonal polynomial L_n has exactly n zeros on $[-1, 1]$ (it is a general property, but it also follows easily from formula (5.1)). Then L'_n has exactly $n-1$ zeros on $[-1, 1]$, and its largest zero is to the left of z_p . Furthermore, L''_n has exactly $n-2$ zeros on $[-1, 1]$, and its largest zero is also to the left of z_p . Since L_n is zero at z_p and one at 1, we conclude that the sign of L''_n on $[z_p, 1)$ is positive, which proves (5.3).

Thus, the function β is strictly increasing on $[z_p, 1)$. $\square$

We continue by defining

$$a(x) := \frac{p}{2} \left(\left(\frac{1+x}{2} \right)^{p-1} + \beta(x) \left(\frac{1-x}{2} \right)^{p-1} \right) L'_n(x)^{-1}, \quad x \in [z_p, 1).$$

Then $(x, a(x), \beta(x)^{1/p})$, $x \in (z_p, 1)$, is a *touching triple*, in the sense that $a(x)L_n$ touches $h_{\beta(x)^{1/p}}$ at the point x . We have already checked that β increases strictly on $[z_p, 1)$ from $\left(\frac{1+z_p}{1-z_p}\right)^p$ to $+\infty$. Let us check now that $a(x)$ decreases strictly on $[z_p, 1)$ from

$$a(z_p) = h'_{\frac{1+z_p}{1-z_p}}(z_p) / L'_n(z_p) \quad (5.4)$$

to 1.

Lemma 14. *The function $a(x)$ is strictly decreasing on $[z_p, 1)$.*

Proof. At the touching point, we have

$$a(x)L_n(x) = h_{\beta(x)^{1/p}}(x).$$

Differentiating both sides gives us

$$a'(x)L_n(x) + a(x)L'_n(x) = h'_{\beta(x)^{1/p}}(x) - \beta'(x) \left(\frac{1-x}{2} \right)^p.$$

Since $h'_{\beta(x)^{1/p}}(x) = a(x)L'_n(x)$, we obtain

$$a'(x)L_n(x) = -\beta'(x) \left(\frac{1-x}{2} \right)^p = -A(x)L_n(x)L''_n(x),$$

where $A(x) > 0$. Since $L_n > 0$ in $(z_p, 1)$ and $L''_n > 0$ in $[z_p, 1)$, the assertion of the lemma follows. $\square$

Summing up, we have unique touching triples $(x, a(x), \beta(x)^{1/p})$ for $x \in [z_p, 1)$ with $a(x) \rightarrow 1$ and $\beta(x) \rightarrow \infty$ as $x \rightarrow 1$. Let $(z_p, a(z_p), \beta(z_p)^{1/p})$ be the triple at z_p . Choose c large enough so that $h_c < a(z_p)L_n$ on the whole interval $[z_p, 1]$. Now decrease c continuously till h_c (h_c increases) first meets $a(z_p)L_n$ at some point $x \in [z_p, 1)$.

If the meeting point x is on $(z_p, 1)$, it must be also a touching point. Then $a(x) = a(z_p)$. On the other hand, $x > z_p$, and so by Lemma 14 one has $a(x) < a(z_p)$. We came to a contradiction. Thus, the meeting point is z_p . Next we verify that it is also a touching point. Indeed, c is equal to $\frac{1+z_p}{1-z_p}$, therefore $h'_c(z_p) = a(z_p)L'_n(z_p)$ by (5.4). This means exactly that h_c (with

this c) not only meets $a(z_p)L_n$ at z_p but also touches it at z_p . Finally, h_c stays below $a(z_p)L_n$ on the whole $(z_p, 1)$: that is how we constructed that c . This completes the proof of Theorem 12.

Here is a different proof of Theorem 12. Let us start with several remarks. First of all, let us fix $1 < a < a(z_p)$ and starting with $c = \infty$ let us decrease c until the first time $h_c(x)$ meets $L_{n,a}(x)$ on $[z_p, 1)$ being below $L_{n,a}(x)$ on the whole interval $[z_p, 1)$. Since $a < a(z_p)$, the meeting points are on $(z_p, 1)$, and hence are touching points.

The touching point on the interval $[z_p, 1)$ (we call it $x(a) = x(n, a)$) is unique. Otherwise, suppose that for some $a > 1$ and for some $c = c(a)$ we have two touching points on $[z_p, 1)$. Then by the previous considerations we have

$$\beta(x_1) = \beta(x_2) = c^p,$$

which contradicts to Lemma 13. We increase a and get decreasing $c(a)$ and touching points $x(a) \in [z_p, 1)$. How can this process end up? The first possibility is that for a certain $a_p > 1$ we have h_c lying below L_{n,a_p} and touching it at z_p . Then automatically $c = c_p := \frac{1+z_p}{1-z_p}$, and $a_p = a(z_p)$; otherwise, the process may stop when $a_p = a(z_p)$. Then again $c = c_p = \frac{1+z_p}{1-z_p}$.

One more proof of Theorem 12 can be obtained from the following fact which we establish for all $p > 2$:

Lemma 15. *The common tangent line ℓ_p separates the graphs of L_{α, a_p} and h_{c_p} on $(z_p, 1)$ if $p = \alpha(\alpha + 1) > 2$.*

Proof. It is easy to calculate the slope of ℓ_p :

$$h'_{c_p}(z_p) = \frac{p}{2} \left(\frac{1+z_p}{2} \right)^{p-1} \left(1 + \frac{1+z_p}{1-z_p} \right) = p \left(\frac{1+z_p}{2} \right)^{p-1} \frac{1}{1-z_p}.$$

Then

$$\ell_p(1) = p \left(\frac{1+z_p}{2} \right)^{p-1}.$$

Later on we prove that

$$p \left(\frac{1+z_p}{2} \right)^{p-1} \geq 1, \quad p \geq 2. \quad (5.5)$$

In Lemma 20 below we prove that L_α is convex on $[z_p, 1]$; hence, L_{α, a_p} lies above ℓ_p . The function h_{c_p} is concave on (z_p, i_p) (recall that i_p denote the unique inflection point of h_{c_p}). So $h_{c_p} < \ell_p$ on $(z_p, i_p]$. In particular, $h_{c_p}(i_p) < \ell_p(i_p)$. On $[i_p, 1]$ the function h_{c_p} is convex. We have seen that at the left end of this interval it is lower than the line ℓ_p . Inequality (5.5) shows that the same happens at the right end point 1, because $h_{c_p}(1) = 1$. This completes the proof of our lemma modulo (5.5).

To verify (5.5) we first establish

Lemma 16. $\frac{1+z_p}{2} \geq \frac{p}{p+2}$, $p \geq 2$.

Proof. We have

$$pL_\alpha(x) = -((1-x^2)L'_\alpha(x))'.$$

We integrate this equality from z_p to 1:

$$(1-z_p^2)L'_\alpha(z_p) = p \int_{z_p}^1 L_\alpha(x) dx.$$

Since L_α is convex on $[z_p, 1]$, the integrand is bigger than $L'_\alpha(z_p)(x - z_p)$. Therefore,

$$(1-z_p^2)L'_\alpha(z_p) \geq \frac{p}{2} L'_\alpha(z_p)(1-z_p)^2.$$

This is equivalent to

$$\frac{1+z_p}{1-z_p} \geq \frac{p}{2}.$$

which proves our assertion. $\square$

To prove (5.5) and to finish the proof of Lemma 15 it remains to mention that

$$\left(\frac{p}{p+2}\right)^{p-1} \geq \frac{1}{p}, \quad p \geq 2,$$

or $p^p \geq (p+2)^{p-1}$. This elementary inequality is true for $p \geq 2$ with equality only for $p = 2$. $\square$

Thus, our best meeting point is z_p , and our best c_p is

$$\frac{1+z_p}{1-z_p}.$$

As $x(p, a) = z_p$ and c_p are already defined, we determine $a = a_p$ from equalities above:

$$a_p = \frac{p}{2} \left(\left(\frac{1+z_p}{2} \right)^{p-1} + \left(\frac{1+z_p}{1-z_p} \right)^p \left(\frac{1-z_p}{2} \right)^{p-1} \right) \cdot L'_n(z_p)^{-1}.$$

5.1. A candidate for the solution of (4.9), (4.10), (4.3). We consider the function (recall that $p = n(n+1)$)

$$g_p(s) := \begin{cases} L_{n,a_p}(s), & s \in [z_p, 1] \\ h_{c_p}(s), & s \in [-1, z_p]. \end{cases}$$

It is C^1 -smooth, and it satisfies (4.9): $Dg_p \leq 0$. In fact, it actually satisfies the equality $Dg_p = 0$ on the interval $(z_p, 1]$. On the interval $[-1, z_p]$, $g_p = h_{c_p}$ (recall that h_{c_p} has an inflection point i_p such that $c_p^p = \left(\frac{1+i_p}{1-i_p} \right)^{p-2}$ located to the right of z_p satisfying the property $c_p^p = \left(\frac{1+z_p}{1-z_p} \right)^p$). Recall also that at the beginning of this section we checked that the inflection point of h_c coincides with the point where Dh_c changes the sign from negative to positive. Therefore, to the left of i_p we have $Dh_{c_p} \leq 0$, which implies that to the left of z_p we have $Dg_p = Dh_{c_p} \leq 0$.

Now let us check (4.10), (4.3). Recall that $\tilde{D}g = \frac{Dg}{1-s^2} + Kg$. Therefore, to check (4.10) (i.e. $\tilde{D}g_p(s) \leq 0$), it is sufficient to prove that

$$-Kg_p(s) = (1-s^2)g_p''(s) + (2p-2)sg_p'(s) - (p^2-p)g_p(s) \geq 0.$$

The function g_p coincides with h_{c_p} on $[-1, z_p]$. Note that

$$Kh_c(s) = 0$$

identically on $[-1, 1]$ for any c . So we need only to check the inequality $-KL_n(s) \geq 0$ on $[z_p, 1]$. We are to verify that

$$(1-s^2)L_n''(s) + (2p-2)sL_n'(s) - (p^2-p)L_n(s) \geq 0,$$

and we have

$$DL_n(s) = (1-s^2)L_n''(s) - 2sL_n'(s) + pL_n(s) = 0. \quad (5.6)$$

Therefore, it suffices to check that

$$p(2sL'_n(s) - pL_n(s)) \geq 0, \quad s \in [z_p, 1].$$

Using (5.6) once more, we see that this follows from (5.3).

Lemma 17. *For $p = n(n+1)$, inequality (4.3) is satisfied for the function $\phi(x, y) = (x+y)^p g_p(\frac{y-x}{x+y})$ such that $g_p = L_{n, a_p}$ to the right of z_p and $g_p = h_{c_p}$ to the left of z_p , where $c_p = \frac{1+z_p}{1-z_p}$.*

Proof. If $s = \frac{y-x}{x+y}$ belongs to $(z_p, 1]$, then $g_p = L_{n, a_p}$, and (4.3) follows, see (4.11). To check (4.3) for others s is easy. Indeed, here $g_p = h_{c_p}$. For $\psi(x, y) := (x+y)^p((y/(x+y))^p - c_p^p(x/(x+y))^p) = y^p - c_p^p x^p$ we have $\psi_{xy} = 0$, $\psi_x/x + \psi_{xx} = -c_p^p p^2 x^{p-2}$, $\psi_y/y + \psi_{yy} = p^2 y^{p-2}$. Inequality (4.3) for ψ can be written as

$$|(\psi_x/x + \psi_{xx})(\psi_y/y + \psi_{yy})|^{1/2} \leq |\psi_x/x + \psi_{xx}|,$$

or, equivalently,

$$y^{p-2} \leq c_p^p x^{p-2}. \quad (5.7)$$

If $s < z_p$, then

$$(1+s)^{p-2} \leq c_p^p (1-s)^{p-2},$$

and (5.7) follows by (4.4). $\square$

Remark 18. We have proved (4.3) for $p = n(n+1)$. However, our argument extends to all $p > 2$.

All the inequalities (4.1), (4.2), (4.3) are now proved for $p = n(n+1)$. This shows that the constant in the orthogonal martingale estimate for such p satisfies the inequality

$$c_p \leq \frac{1+z_p}{1-z_p}.$$

6. LEGENDRE EQUATION. GENERAL FACTS

Here we list some useful facts from [14].

Consider the equation

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

near the point $x = x_0$, where P, Q, R are analytic functions and

$$(x - x_0) \frac{Q(x)}{P(x)}, \quad (x - x_0)^2 \frac{R(x)}{P(x)}$$

are analytic functions in a neighborhood of x_0 (the regular singular point case). Suppose that the *indicial* polynomial

$$r(r - 1) + a_0r + b_0,$$

where

$$a_0 = \lim_{x \rightarrow x_0} (x - x_0) \frac{Q(x)}{P(x)}, \quad b_0 = \lim_{x \rightarrow x_0} (x - x_0)^2 \frac{R(x)}{P(x)}$$

has a double root $r = r_1$. Then our equation has two linearly independent solutions represented in a small half-neighborhood $(x_0, x_0 + \varepsilon)$ by the formulas

$$f_1(x) = (x - x_0)^{r_1} \sum_{n=0}^{\infty} a_n (x - x_0)^n,$$

$$f_2(x) = f_1(x) \log \frac{1}{x - x_0} + (x - x_0)^{r_1} \sum_{n=1}^{\infty} b_n (x - x_0)^n;$$

the series converge absolutely for $x \in (x_0 - \varepsilon, x_0 + \varepsilon)$, $a_0 \neq 0$.

In the case of the Legendre equation,

$$(1 - x^2)y'' - 2xy' + py = 0, \tag{6.1}$$

we will use these notations for a bounded and an unbounded solutions near $x_0 = 1$:

$$f_1(x) = 1 + \frac{p}{2 \cdot 1^2}(x - 1) - \frac{p(1 \cdot 2 - p)}{(2 \cdot 1^2)(2 \cdot 2^2)}(x - 1)^2 + \dots$$

$$+ (-1)^{n+1} \frac{p(1 \cdot 2 - p) \dots (n(n - 1) - p)}{2^n (n!)^2} (x - 1)^n + \dots$$

For integer α this is a Legendre polynomial. Furthermore, in a half-neighborhood $(1 - \varepsilon, 1)$ we have

$$f_2(x) = f_1(x) \log \frac{1}{1-x} + H(x),$$

where H is real analytic in a neighborhood of 1; f_1 and f_2 are real analytic on $(-1, 1)$.

7. THE CASE $p > 2$

To extend our solution of the main problem from the case $p = n(n+1)$ to the general case $p > 2$, we need to prove a couple of lemmas. Denote by z_p the rightmost zero of f_1 on the interval $[-1, 1]$. We are going to prove two things: (1) for every solution of Legendre equation (6.1), its rightmost zero on the interval $[-1, 1]$ is at least z_p ; (2) the solution f_1 is strictly convex on $[z_p, 1]$.

We need also Remark 18 to complete the reasoning for all $p > 2$.

Lemma 19. *For every solution of Legendre equation (6.1), its rightmost zero on the interval $[-1, 1]$ is at least z_p .*

Proof. Note that $f_2(1) = +\infty$. Consider the Wronskian $W(x) = f_2'(x)f_1(x) - f_1'(x)f_2(x)$. Section 6 gives us that $W(x) \asymp \frac{1}{1-x}$, $x < 1$. So W is positive near $x = 1$. We know that

$$W'(x) = \frac{2x}{1-x^2}W(x),$$

and, hence, W preserves the sign. Consider

$$W(z_p) = f_2'(z_p)f_1(z_p) - f_1'(z_p)f_2(z_p) = -f_1'(z_p)f_2(z_p).$$

The function f_1 is positive on $[z_p, 1]$, and it changes sign at z_p , so $f_1'(z_p) \geq 0$. If $f_1'(z_p) = 0$, then by the Legendre equation, $f_1''(z_p) = 0$, and differentiating the Legendre equation, we get $f_1^{(n)}(z_p) = 0$ for all n . This is impossible as the analytic function f_1 would then vanish identically. Hence, $f_1'(z_p) > 0$. We conclude that

$$f_2(z_p) < 0.$$

Consider a linear combination $f_3 = c_1 f_1 + c_2 f_2$ with a positive c_2 . Then $f_3(z_p) < 0$ by what we have just proved. Since $f_3(1) = +\infty$, f_3 must have a zero on $(z_p, 1)$. For negative c_2 we have the same conclusion. $\square$

Lemma 20. *The function f_1 is strictly convex on $[z_p, 1)$ for $p > 2$.*

Proof. We have

$$\begin{aligned} (1 - x^2)f_1'' - 2xf_1' + pf_1 &= 0, \\ (1 - x^2)f_1''' - 4xf_1'' + (p - 2)f_1' &= 0. \end{aligned}$$

Then

$$f_1'(1) = \frac{p}{2}, \quad f_1''(1) = \frac{p(p-2)}{8}.$$

We have already observed that $f_1'(z_p) > 0$. By the first equation above we get $f_1''(z_p) > 0$.

Let $x_1 \in (z_p, 1)$ be the first point where $f_1''(x_1) = 0$. Then obviously $f_1'(x_1) > 0$, and by the second equation above $f_1'''(x_1) < 0$. So f_1 does change convexity to concavity passing through x_1 . Furthermore, we have just seen that $f_1''(1) = \frac{p(p-2)}{8} > 0$. So f_1 should change from concavity to convexity again, say at $x_3 \in (x_1, 1)$. Let x_3 be the closest to x_1 point with this property, so that $f_1'' < 0$ in between. Then there exists $x_2 \in (x_1, x_3)$ such that $f_1'''(x_2) = 0$, $f_1''(x_2) < 0$. Plug this to the second equation at the beginning of the proof and note that then

$$f_1'(x_2) < 0.$$

Therefore, using again the Legendre equation we see that f_1', f_1'' will stay negative till the point 1. This is impossible because they are strictly positive at 1. $\square$

Let us observe that Remark 18 and Lemmas 19, 20 are the only ingredients we need to carry through the reasoning for general $p > 2$. We just repeat the reasoning we used for the case $p = n(n+1)$ replacing the Legendre polynomials L_n by f_1 . We finally get the proof that for $p > 2$ the best constant for the

martingale transform of orthogonal martingales satisfies the inequality

$$c_p \leq \frac{1 + z_p}{1 - z_p},$$

where z_p is the largest zero of the solution f_1 of Legendre equation (6.1): $(1 - x^2)y'' - 2xy' + py = 0$ on the interval $[-1, 1]$. We also proved that $z_p = \min_y \max\{z(y)\}$, where $z(y)$ denotes any zero z of any nontrivial solution y of (6.1) on the interval $[-1, 1]$.

8. SHARPNESS

We use the spherical coordinates

$$x = r \sin(\theta) \cos(\phi), y = r \sin(\theta) \sin(\phi), z = r \cos(\theta).$$

The $\mathbb{R}^3$ Laplacian in spherical coordinates is

$$\Delta\psi = \frac{\partial_r(r^2\partial_r\psi)}{r^2} + \frac{\partial_\theta(\sin(\theta)\partial_\theta\psi)}{r^2\sin\theta} + \frac{\partial_\phi^2\psi}{r^2\sin^2\theta}.$$

Changing the variable $s = \cos\theta$, we define the obstacle function

$$v(\theta) = \left(\frac{1 + \cos\theta}{2}\right)^{\alpha(\alpha+1)} - c^{\alpha(\alpha+1)} \left(\frac{1 - \cos\theta}{2}\right)^{\alpha(\alpha+1)}.$$

We associate to v an auxiliary obstacle function in $\mathbb{R}^3$,

$$V(r, \theta, \phi) = r^\alpha v(\theta).$$

Observe that the function V has separated variables and azimuthal symmetry (i.e. no dependence on ϕ).

Consider now the *minimal* superharmonic function ψ , $\psi \geq V$. Then ψ has the same symmetries as V does. This is a consequence of taking infimum over the superharmonic majorants of the form $\psi(ax + by, -bx + ay, z)$ and $\frac{1}{\lambda^\alpha}\psi(\lambda x, \lambda y, \lambda z)$; this infimum which has both homogeneity and rotational invariance is again superharmonic. Thus the minimal superharmonic majorant $\psi(r, \theta, \phi) \geq V$ has the form

$$\psi(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi) = r^\alpha\Theta(\theta).$$

Applying the spherical Laplacian and using the azimuthal symmetry, we obtain (at least in the sense of distributions) that

$$r^2 \sin(\theta) \Delta \psi = \alpha(\alpha + 1) r^\alpha \Theta(\theta) \sin \theta + r^\alpha (\Theta'(\theta) \sin \theta)'.$$

Dividing by $r^\alpha \sin \theta$, we get

$$\frac{\Delta \psi}{r^{\alpha-2}} = \frac{(\Theta'(\theta) \sin \theta)'}{\sin \theta} + \alpha(\alpha + 1) \Theta(\theta).$$

Here $\theta \in (0, \pi)$. The right hand side is the trigonometric form of the Legendre operator. The usual form $((1 - s^2)y')' + \alpha(\alpha + 1)y$ is obtained from this by the substitution $s = \cos \theta$ and $y(\cos \theta) = \Theta(\theta)$. In particular, we obtain that seeking for the minimal $y(s) \geq v(s) = h_c(s)$ such that

$$((1 - s^2)y')' + \alpha(\alpha + 1)y \leq 0$$

is equivalent to seeking for the minimal superharmonic $\psi \geq V$. It is well-known that ψ is harmonic wherever $\psi > V$. We know that the only harmonic function satisfying these homogeneity and symmetry conditions corresponds to a solution of the Legendre equation.

We recall that the Bellman function $\mathcal{B}$ from the beginning of the paper generates a function g on $[-1, 1]$ satisfying (4.9), (4.10). We use only (4.9), which shows that this particular g is a supersolution of the Legendre equation: $(1 - x^2)g'' - 2xg' + pg \leq 0$. Thus, the function g generates a superharmonic majorant of V given by the formula

$$\Psi(r, \theta, \phi) = r^\alpha g(\cos(\theta)).$$

Since $\psi \leq \Psi$, there exists another solution of (4.9) which actually satisfies the Legendre equation everywhere, where it is strictly bigger than h_c . Denote this solution of the Legendre equation by L . When L and h_c meet at a certain point s_0 , it should be a point where they are tangent to each other. Otherwise, if $L > h_c$ on $(s_0, s_0 + \epsilon)$, and $L(s_0) = h_c(s_0)$, then $L'(s_0) > h'_c(s_0)$. Define a new function: $f = L$ to the right of s_0 and $f = h_c$ to the left of s_0 . It cannot be a solution of (4.9). In fact, the inequality $L'(s_0) > h'_c(s_0)$ implies that $f''(s_0)$ is a positive delta function, and so satisfies at s_0 the inequality exactly opposite to (4.9).

Thus, L is inevitably tangent to h_c at a meeting point. Furthermore, such pairs (L, h_c) , where L is a solution of the Legendre equation and h_c is an obstacle function, were already considered when we were looking for the smallest constant c . Thus, the minimal possible c is $\frac{1+z_p}{1-z_p}$.

9. ANOTHER WAY TO PROVE SHARPNESS

The use of the Laplacian in $\mathbb{R}^3$ is of course very specific for the Legendre equation. The Legendre equation and spherical harmonics are close relatives. We want to show an approach that uses much less specifics of the ODE (still, it uses some of it).

Let

$$P(x)y'' + Q(x)y' + R(x)y = 0$$

be an equation such that the functions P, Q, R are real analytic, and let x_0 be a regular singular point (we choose $x_0 = 1$ as in the Legendre case, but this is of no importance). One has two solutions: a bounded one, f_1 , and an unbounded one, f_2 . In particular, this happens when the indicial equation has a double root $r_1 = 0$ as in the Legendre case. The reader may take a look at Section 6. We assume here that $f_1(1) > 0$ and (unlike in Section 6) $f_2(1) = -\infty$.

Let b be the closest to 1 zero of f_1 , $b < 1$. We deal only with the interval $[b, 1]$, and assume that P, Q, R keep sign on $[b, 1)$.

Lemma 21. *Let g be a supersolution on $[b, 1]$, that is $P(x)g'' + Q(x)g' + R(x)g \leq 0$, and let $g(b) > 0$. Then $g(1) = -\infty$.*

Let us first explain why this lemma implies the sharpness of the constant c_p . Recall that the Bellman function $\mathcal{B}$ from the beginning of the paper generates a function g on $[-1, 1]$ satisfying (4.9), (4.10). We use only (4.9), which shows that this particular g is a supersolution of the Legendre equation: $(1 - x^2)g'' - 2xg' + pg \leq 0$. We constructed a special solution L_{p,a_p} with the “best” zero z_p , positive on $(z_p, 1]$. The obstacle function h_{c_p} lies below L_{p,a_p} and touches it at z_p . If our g were such that $g \geq h_c$ on $[-1, 1]$, and $c < c_p$, then $g \geq h_c(z_p) > 0$. Lemma 21 shows that the inequality $g \geq h_c$ is impossible on

$[z_p, 1]$ because $h_c(1) = 1$, $g(1) = -\infty$. Therefore, we must have $c \geq c_p$. And this is exactly the sharpness of our constant.

It remains to prove our lemma.

Proof of Lemma 21. Consider two Wronskians:

$$\begin{aligned} W(x) &= f_2'(x)f_1(x) - f_1'(x)f_2(x), \\ \tilde{W}(x) &= g'(x)f_1(x) - f_1'(x)g(x). \end{aligned}$$

Using that g is a supersolution, and f_1 is a solution we can write on $(b, 1)$:

$$\tilde{W}' \leq -\frac{Q}{P}\tilde{W}.$$

On the other hand, we have everywhere

$$W' = -\frac{Q}{P}W.$$

Combining these two relations we get

$$\tilde{W}' \leq \frac{W'}{W}\tilde{W}. \quad (9.1)$$

Furthermore, it is easy to see that $\lim_{x \rightarrow 1} W(x) = -\infty$ (because of the behavior of f_2). The Wronskian W preserves the sign, so $W(x) < 0$ on $[b, 1]$. Therefore, (9.1) (after division by $W < 0$) can be rewritten as

$$(\tilde{W}/W)' \geq 0 \text{ on } [b, 1]. \quad (9.2)$$

Note that $\tilde{W}(b) = -g(b)f_1'(b) < 0$, $W(b) = -f_2(b)f_1'(b) < 0$. Set $\kappa = \frac{g(b)}{f_2(b)} > 0$. Inequality (9.2) shows that

$$\tilde{W}(x)/W(x) \geq \kappa$$

on $[b, 1]$. Using the negativity of W , we get

$$\tilde{W}(x) \leq \kappa W(x), \quad x \in [b, 1]. \quad (9.3)$$

Note that $(g/f_1)' = \tilde{W}/f_1^2$, $(f_2/f_1)' = W/f_1^2$. Then (9.3) becomes

$$\left(\frac{g}{f_1}\right)' \leq \kappa \left(\frac{f_2}{f_1}\right)' \text{ on } [b, 1].$$

Hence,

$$\frac{g}{f_1} \leq \kappa \frac{f_2}{f_1} + \text{const on } [b, 1].$$

Now we see that $\lim_{x \rightarrow 1} g(x) = -\infty$. The lemma is proved. $\square$

10. ASYMPTOTICS OF z_p

The Mehler-Heine formula (1868),

$$\lim_{n \rightarrow \infty, n \in \mathbb{N}} L_n\left(\cos \frac{x}{n}\right) = J_0(x)$$

establishes a relation between the Legendre polynomials L_n and the Bessel function of zero order J_0 . As a consequence ([34, Theorem 8.1.2]),

$$\lim_{n \rightarrow \infty, n \in \mathbb{N}} n(n+1)(1 - z_{n(n+1)}) = \frac{j_0^2}{2}, \quad (10.1)$$

where j_0 is the first positive zero of the Bessel function of zero order J_0 .

Let $\beta > \alpha > 1$, $D_\alpha f_\alpha = 0$, $D_\beta f_\beta = 0$, $f_\alpha(1) = f_\beta(1) = 1$, and let $f_\alpha(x) > 0$, $x_\alpha < x \leq 1$, $f_\alpha(x_\alpha) = 0$. If $f_\beta(x) > 0$ on $[x_\alpha, 1]$, then $D_\alpha f_\beta \leq 0$ on $[x_\alpha, 1]$, and, hence, f_β is a positive supersolution for D_α on $[x_\alpha, 1]$ which is impossible by Lemma 21. Thus, z_p increases for $p \in (2, +\infty)$, and (10.1) gives us

$$\lim_{p \rightarrow \infty} p(1 - z_p) = \frac{j_0^2}{2}.$$

11. ON BURKHOLDER FUNCTIONS

We did not find the formula for the function $\mathcal{B}$ from Section 3.1. However, we have found the formula for the function $\Phi = \Phi_{c_p}$ from Section 3.2. Indeed, the function ϕ of two variables associated to Φ_{c_p} gives rise to a function g_p of one variable on $[-1, 1]$ such that the corresponding function $r^\alpha g_p(\cos \theta)$ (recall that $p = \alpha(\alpha + 1)$) expressed in the spherical coordinates in $\mathbb{R}^3$ is the least superharmonic majorant of

$$V(r, \theta, \phi) := r^\alpha \left(\left(\frac{1 + \cos \theta}{2} \right)^p - c_p^p \left(\frac{1 - \cos \theta}{2} \right)^p \right).$$

How to recognize the least superharmonic majorant of a given function V ? This is the function which is harmonic everywhere where it is strictly bigger than V . We constructed (using our g_p) exactly such a function. So we can completely restore Φ from g_p .

12. ONE-SIDED ORTHOGONALITY. ANSWERS AND QUESTIONS.

Suppose that Z is orthogonal, and W is not. We always assume that W is subordinate to Z . For $1 < p \leq 2$ we have cited an estimate

$$\|W\|_p \leq \sqrt{\frac{2}{p(p-1)}} \|Z\|_p.$$

Is this sharp? We do not know.

What if $p \geq 2$? Here is a result which we prove in [6]:

$$\|W\|_p \leq \sqrt{2} \frac{1 - s_p^*}{s_p^*} \|Z\|_p, \quad p \geq 2,$$

where s_p^* is the closest to 0 zero of a bounded near 0 solution of the Laguerre equation

$$s^2 \mathcal{L}'' + (1 - s) \mathcal{L}' + p \mathcal{L} = 0. \quad (12.1)$$

This estimate is sharp, see [6].

Suppose that W is orthogonal, and Z is not. For $p \geq 2$ we have cited an estimate

$$\|W\|_p \leq \sqrt{\frac{p(p-1)}{2}} \|Z\|_p.$$

Is this sharp? We do not know.

What if $1 < p \leq 2$? Here is a result which we prove in [6]:

$$\|W\|_p \leq \frac{1}{\sqrt{2}} \frac{s_p}{1 - s_p} \|Z\|_p, \quad 1 < p \leq 2,$$

where s_p is the closest to 1 zero of a bounded near 0 solution of Laguerre equation (12.1). This estimate is sharp, see [6].

REFERENCES

- [1] A. BAERNSTEIN AND S. MONTGOMERY-SMITH, *Some conjectures about integral means of ∂f and $\bar{\partial} f$* , Complex analysis and Differential Equations, Proc. of the Marcus Wallenberg symposium in honor of Matts Essén (Uppsala, Sweden, 1997), 92–109, 1999.
- [2] R. BAÑUELOS, P. JANAKIRAMAN, *L^p -bounds for the Beurling–Ahlfors transform*. Trans. Amer. Math. Soc. **360** (2008), 3603–3612.
- [3] R. BAÑUELOS, P. J. MÉNDEZ-HERNÁNDEZ, *Space-time Brownian motion and the Beurling–Ahlfors transform*, Indiana Univ. Math. J. **52** (2003), 981–990.

- [4] R. BAÑUELOS, A. OSEKOWSKI, *Burkholder inequalities for submartingales, Bessel processes and conformal martingales*, Preprint, 2011, pp. 1–20.
- [5] R. BAÑUELOS, G. WANG, *Sharp inequalities for martingales with applications to the Beurling–Ahlfors and Riesz transforms*, Duke Math. J. **80** (1995), 575–600.
- [6] A. BORICHEV, P. JANAKIRAMAN, A. VOLBERG, *Subordination by orthogonal martingales in L^p and zeros of Laguerre polynomials*, Preprint, 2010, 30 pp.
- [7] D. BURKHOLDER, *Boundary value problems and sharp estimates for the martingale transforms*, Ann. of Prob. **12** (1984), 647–702.
- [8] D. BURKHOLDER, *An extension of classical martingale inequality*, Probability Theory and Harmonic Analysis, ed. by J.-A. Chao and W. A. Woyczynski, Marcel Dekker, 1986.
- [9] D. BURKHOLDER, *Sharp inequalities for martingales and stochastic integrals*, Colloque Paul Lévy sur les Processus Stochastiques (Palaiseau, 1987), Astérisque No. 157–158 (1988), 75–94.
- [10] D. BURKHOLDER, *A proof of the Peczyński’s conjecture for the Haar system*, Studia Math. **91** (1988), 79–83.
- [11] D. BURKHOLDER, *Differential subordination of harmonic functions and martingales*, (El Escorial 1987) Lecture Notes in Math. **1384** (1989), 1–23.
- [12] D. BURKHOLDER, *Explorations of martingale theory and its applications*, Lecture Notes in Math. **1464** (1991), 1–66.
- [13] D. BURKHOLDER, *Strong differential subordination and stochastic integration*, Ann. of Prob. **22** (1994), 995–1025.
- [14] E. A. CODDINGTON, *An Introduction to Ordinary Differential Operators*. New York, Dover, 1989.
- [15] O. DRAGICEVIC, A. VOLBERG *Sharp estimate of the Ahlfors–Beurling operator via averaging martingale transforms*, Michigan Math. J. **51** (2003), 415–435.
- [16] O. DRAGICEVIC, A. VOLBERG, *Bellman function, Littlewood–Paley estimates, and asymptotics for the Ahlfors–Beurling operator in $L^p(\mathbb{C})$* , Indiana Univ. Math. J. **54** (2005), 971–995.
- [17] O. DRAGICEVIC, A. VOLBERG, *Bellman function and dimensionless estimates of Littlewood–Paley type*, J. of Oper. Theory **56** (2006) 167–198.
- [18] O. DRAGICEVIC, S. TREIL, A. VOLBERG, *A lemma about three quadratic forms*, Intern. Math. Research Notices. 2008, 9 pp.
- [19] S. GEISS, S. MONTGOMERY-SMITH, E. SAKSMAN, *On singular integral and martingale transforms*, Trans. Amer. Math. Soc. **362** (2010), 553–575.
- [20] I. KARATZAS, S. SHREVE, *Brownian Motion and Stochastic Calculus*. Graduate Texts in Mathematics, Springer, 1991.

- [21] N. KRYLOV, *Controlled Diffusion Processes*, Springer, 1980.
- [22] F. NAZAROV, S. TREIL, *The hunt for Bellman function: applications to estimates of singular integral operators and to other classical problems in harmonic analysis*, St. Petersburg Math. J. **8** (1997), 721–824.
- [23] F. NAZAROV, S. TREIL, A. VOLBERG, *The Bellman functions and two-weight inequalities for Haar multipliers*, J. Amer. Math. Soc. **12** (1999), 909–928.
- [24] F. NAZAROV, S. TREIL, A. VOLBERG, *Bellman function in stochastic control and harmonic analysis*, Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), 393–423, Oper. Theory Adv. Appl., 129, Birkhauser, Basel, 2001.
- [25] F. NAZAROV AND A. VOLBERG, *The Bellman function, two-weight Hilbert transform, and embeddings of the model spaces K_θ* , Dedicated to the memory of Thomas H. Wolff, J. Anal. Math. **87** (2002), 385–414.
- [26] F. NAZAROV AND A. VOLBERG, *Heating of the Ahlfors–Beurling operator and estimates of its norm*, St. Petersburg Math. J. **15** (2004), 563–573.
- [27] S. PETERMICHL, *The sharp bound for the Hilbert transform on weighted Lebesgue spaces in terms of the classical A_p characteristic*, Amer. J. Math. **129** (2007), 1355–1375.
- [28] S. PETERMICHL, A. VOLBERG, *Heating of the Ahlfors–Beurling operator: weakly quasiregular maps on the plane are quasiregular*, Duke Math. J. **112** (2002), 281–305.
- [29] S. PICHORIDES, *On the best values of the constants in the theorems of M. Riesz, Zygmund, and Kolmogorov*, Studia Math. **44** (1972) 165–179.
- [30] A. V. POGORELOV, *Extrinsic geometry of convex surfaces*. Translations of Mathematical Monographs, Amer. Math. Soc., v. 35, 1973.
- [31] E. SAWYER, *Two weight norm inequalities for certain maximal and integral operators*, Lecture Notes Math., **908**, Springer-Verlag, Berlin–Heidelberg–New York, 1982, pp. 102–127.
- [32] L. SLAVIN, A. STOKOLOS, *The Bellman PDE for the dyadic maximal function and its solution*, Preprint, 2006.
- [33] L. SLAVIN, V. VASYUNIN, *Sharp results in the integral-form John–Nirenberg inequality*, Trans. AMS **363** (2011) 4135–4169.
- [34] G. SZEGÖ, *Orthogonal polynomials*, Fourth edition. American Mathematical Society, Colloquium Publications, Vol. XXIII, American Mathematical Society, Providence, R.I., 1975.
- [35] V. VASYUNIN, A. VOLBERG, *The Bellman function for a certain two-weight inequality: a case study*, St. Petersburg Math. J. **18** (2007), 201–222.

- [36] V. VASYUNIN, A. VOLBERG, *Monge–Ampère equation and Bellman optimization of Carleson Embedding Theorems*, arXiv:0803.2247. Advances in Math. Sciences, Ser. 2, v. 226, (2009), pp. 195–238. Amer. Math. Soc. Translations.
- [37] V. VASYUNIN, A. VOLBERG, *Bellman Functions Technique in Harmonic Analysis*, preprint (sashavolberg.wordpress.com), 2009, 86 pp.
- [38] A. VOLBERG, *Bellman approach to some problems in Harmonic Analysis*, Séminaires des Equations aux dérivées partielles. Ecole Polytechnique, 2002, exposé XX, pp. 1–14.
- [39] G. N. WATSON, *A treatise on the theory of Bessel functions*, Reprint of the second (1944) edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1995.

ALEXANDER BORICHEV,
UNIVERSITÉ DE PROVENCE, MARSEILLE,
borichev@cmi.univ-mrs.fr,

PRABHU JANAKIRAMAN,
DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY,
pjanakir1978@gmail.com,

ALEXANDER VOLBERG,
DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY,
volberg@math.msu.edu